Keys Please! A Guide to Classification



Specific Learning Outcomes 6-1-01: Use appropriate vocabulary related to their investigations of the diversity of living things.

6-1-02: Describe various kinds of classification systems used in everyday life, and identify related advantages and disadvantages.
6-1-04: Identify living things using an existing classification key, and explain the rationale used.
6-1-05: Identify advantages and disadvantages of having a common classification system for living things, and recognize that the system changes as new evidence comes to light.

General Learning Outcomes

6-0-1a: Formulate specific questions that lead to investigations.

6-0-1b: Identify various methods for finding the answer to a specific question and select one to implement.

6-0-2a: Access information using a variety of sources.

6-0-2b: Review information to determine its usefulness, using predetermined criteria.

6-0-2c: Make notes on a topic, combing information from more than one source and referencing sources appropriately.

6-0-5a: Make observations that are relevant to a specific question.

6-0-6f: Evaluate the methods used to answer a question or solve a problem.

Vocabulary

classification system, classification key, shorebird, wetland

Summary

Students are introduced to wetlands and the diversity of living things by developing the ability to use a classification key in order to identify shorebirds found at Oak Hammock Marsh. Students will also evaluate the usefulness of using a key, as well as the advantages and disadvantages of having a common classification system for living things.

Materials

- Projector and computer to present slideshow
- Student access to a library and the internet
- Print one key per student
- Either print images in colour and cut-out (recommend laminating for future use) or provide a digital copy of the shorebird pictures and identification tips page for student assignment

Procedure

Warm Up

Begin with the provided slideshow presentation, which discusses the need for classification when studying living things and introduces students to different types of classification systems. The slideshow will be both the warm up and the majority of the activity.

A wetland is an area of land that holds shallow water, with a maximum depth of two metres. The water makes the soil very moist, so that water-loving plants will grow in and around the wetland; this is why a wetland can not be deeper then two metres, because otherwise these kinds of plants drown and do not receive enough sunlight. The water moves slowly because there are so many plants that slow the flow, absorbing some of the water like a sponge and filtering it as it moves through.

A classification system is an approach to classification, where a large group is divided into smaller subgroups then arranged in a particular way, used to better differentiate between those in a group.

A classification key is a tool that helps determine the identity of something in the natural world through the process of elimination. The key gives you choices based on structural or behavioural features, and as you select one it narrows down the possibilities until you get to a species. If you have correctly used the key and have chosen the right kind of key(s), you arrive at the correct name of the species you are trying to identify.

Activity

The slideshow will provide an opportunity for students to learn how to use a classification key, going through an example step by step.

Each student should be given a copy of the Shorebird classification key which is used in the slideshow as the example, and will also be used for the second half of this activity.

Once you have finished presenting the slideshow, either assign or have students choose one of the shorebird pictures (provided). Each student will then have to go through the key on their own and see if they are able to identify the shorebird they have picked. This can either be a classroom activity (as long as your students have access to the internet) or a take-home assignment.

<u>Recommended</u>: Encourage students to reference the provided Shorebird Identification Tips page, as well as a bird book or the following websites to compare and contrast between the likely possibilities.

Recommended Websites:

- www.allaboutbirds.org
- www.audubon.org

Wrap Up

After students have had the opportunity to identify their shorebird, have students come together to discuss first what bird species they believe they have picked, and than what was their experience with the key.

Conclude by explaining that as a class you will be visiting Oak Hammock Marsh, which is one of the many wetland habitats these shorebirds visit as they move from place to place. Students will have an opportunity to explore and hopefully see a few shorebirds themselves!

<u>Optional:</u> Bring the shorebird classification key or use the Merlin app (which is a free, downloadable bird identification tool) while on your field trip to gain experience using these classification tools in the field! You will see the most shorebirds at Oak Hammock Marsh during the early spring and early fall seasons, although some stay here for the summer to breed.


Identification Tips:

When students are looking at their Shorebird picture, encourage them to notice...

- Body size
- Leg colour
- Bill colour, shape, and length in comparison to their head
- Colour of the plumage (feathers) on their back
- Colour and plumage pattern on the chest - Are there any streaks (vertical lines) or bars (horizontal lines)? How far along the body do the streaks or bars go?
- Colour and patterns around their eyes — Is there an eye ring? Is there an eye line?
- Are the wing feathers longer then the tail?

See accompanying Identification Tips page for visual examples.






Key Characteristics of Bogs:

- Peat-covered (peat is a brown, soil-like substance made of decaying Sphagnum mosses)
- Isolated from ground water
- Rain-fed
- Low nutrients in the water and acidic
- Dominated by Sphagnum mosses with tree, shrub or treeless vegetation cover •

Wetland—Fen



Key Characteristics of Fens:

- Peat-covered (peat is a brown, soil-like substance made of decaying Sphagnum mosses)
- Exposed to ground water; water at surface is moving
- Fed by rain, streams and groundwater
- More nutrients in the water then bogs and is less acidic
- Greater variety of plants than bogs: grass meadows, shrubs, and trees

Wetland—Swamp




Key Characteristics of Swamps:

- Non-peat forming wetland
- Has flowing water; flooded for the majority of the •

growing season

- Waterlogged soil, often standing water
- Vegetation is dense, and can include coniferous or deciduous trees, or tall shrub thickets


Wetland—Freshwater Marsh



Key Characteristics of Freshwater Marshes:

- Non-peat forming wetland
- Frequently or continually flooded with shallow, slow moving water •
- Waterlogged soil that is oxygenated, which allows for plants to form roots •
- Nutrient rich water offers greater plant diversity, such as cattails, reeds, rushes, or sedges
- No trees

Shallow Open Water



Key Characteristics of Shallow Open Water Wetlands:

- Locally known as ponds, sloughs and marshes
- Non-peat forming wetland
- Standing water, often a transition between lake and marsh
- Fewer emergent plants but submersed plants may be present



Shorebird Key

(Adapted from: A key for breeding or partial breeding plumage shorebirds in Prairie Canada – prepared by The Western Hemisphere Shorebird Reserve Network)

This key is designed to help you identify shorebirds found here at Oak Hammock Marsh. It gives you choices and as you select one it narrows down the possibilities until you get to one. This should be the correct species. It is always a good idea to check a book once you have identified your bird. You may still not be satisfied that the identification is right. You can run through the key again and see if you come up with the same answer. It is always possible that the bird you are looking at is not in the key. If this is the case it is probably a rare shorebird.

- I) Body size larger than a Mourning Dove and body colours mostly black and white Avocets
 - A) Bill upturned, legs bluish-grey, peachy coloured head and neck AMERICAN AVOCET
- II) Large head, large eye, short neck, short and stout bill; typically forages in a run/stop & look/run & grab manner

<u>Plovers</u>

- A) Body larger than a robin
 - 1) Plumage is black on upper & lower breast, under-tail is white BLACK-BELLIED PLOVER
 - 2) Plumage is black on upper & lower breast, under-tail is black AMERICAN GOLDEN-PLOVER
 - 3) 2 distinct black bands across white breast KILLDEER
- B) Body smaller than a robin
 - Back plumage light grey or grey-brown, legs light, bill orange with black tip, 1 distinct black breast band
 PIPING PLOVER
 - 2) Back plumage solid brown, 2 distinct black breast bands KILLDEER
 - 3) Back plumage solid brown, 1 distinct black breast band —SEMIPALMATED PLOVER
- III) Small head, small eye, thin bill and neck, forages by probing Sandpipers
 - A) Body equal to or larger than a Mourning Dove
 - 1) Upland habitat, bill same length as head, plumage brown, sits on fence posts UPLAND SANDPIPER
 - 2) Wetland habitat, bill longer than head, long legs
 - (a) Legs greyish, bill straight and thick, black & white wing pattern WILLET
 - (b) Legs yellow
 - (i) Bill less than 1.25X the head length LESSER YELLOWLEGS
 - (ii) Bill greater than 1.25X the head length GREATER YELLOWLEGS







- (c) Legs dark, bill curves up
 - (i) Brown plumage, mottled back, brown barred front MARBLED GODWIT
 - (ii) Black & white tail pattern, reddish front HUDSONIAN GODWIT
- B) Body smaller than a Mourning Dove and larger than a Red-winged Blackbird
 - 1) Bill 2X length of head
 - (a) Typically damp, second-growth woodland, chunky appearance, **usually solitary** AMERICAN WOODCOCK
 - (b) Lateral head stripe, typically wet, grassy habitats, usually solitary or in loose flocks WILSON'S SNIPE
 - (c) Open, muddy habitats, steady probing (sewing-machine-like), usually in closely spaced flocks
 - (i) **Tail looks white with black bars**, white on belly, white rump patch far up back SHORT-BILLED DOWITCHER
 - (ii) Tail looks black with white bars, rusty lower belly, barred sides LONG-BILLED DOWITCHER
 - 2) Bill less than 1.5X length of head
 - (a) Thin, needle-like bill, small head, frequently swimming (often in circles)
 - (i) Legs yellowish, bill longer than head WILSON'S PHALAROPE
 - (ii) Legs dark, bill about length of head RED-NECKED PHALAROPE
 - (b) Long legs, upright posture, conspicuous eye ring
 - (i) **Chronic teetering motion**, smoky brown back, bill length about same as head length, when flying shows white wing stripe SPOTTED SANDPIPER
 - (ii) Occasional teetering motion, fresh water habitat, dark brown back, bill longer than head, when flying shows no white wing stripe — SOLITARY SANDPIPER
 - (iii) Legs yellow, bill less than 1.25X length of head LESSER YELLOWLEGS
 - (c) Long legs, upright posture, no eye ring
 - (i) Legs straw-coloured, **buffy coloration above and below**, typically in short-grass or stubble fields,

long wings — BUFF-BREASTED SANDPIPER

- (d) Hunched posture, eye ring not distinct
 - (i) Straw-coloured legs, brown plumage with **vertical streaks** on upper breast stopping mid-breast in **abrupt line** PECTORAL SANDPIPER
 - (ii) Dark legs, no vertical streaks, **salmon-coloured breast/belly in breeding plumage** RED KNOT







- (iii) Greenish legs, bill slightly down-turned at tip, prominent eye line, typically wading (often bellydeep) and probing for bottom-dwelling invertebrates, often submerging entire head —STILT SANDPIPER
- 3) Bill chisel-shaped, legs orange, conspicuous black chest markings (not bands) RUDDY TURNSTONE
- C) Body smaller than a Red-winged Blackbird
 - 1) Wing tips extend 2+ mm beyond tail
 - (a) **Prominent eye line**, speckles on sides, grey-brown back, typically wading WHITE-RUMPED SANDPIPER
 - (b) Thin, tapered bill, buffy breast with streaks, **brown back**, dark legs, often forages above water line BAIRD'S SANPIPER
 - 2) Wing tips **do not** extend much beyond tail
 - (a) Body slightly larger than a sparrow
 - (i) Clear white belly feathers, light grey back, typically running between foraging attempts
 SANDERLING
 - (ii) Dirty white belly feathers (black belly with reddish back feathering in breeding plumage),
 typically walks between foraging attempts, bill slightly down-curved entire length DUNLIN
 - (b) Body similar in size to a sparrow
 - (i) Smudged and streaked upper breast, brown back, yellowish legs LEAST SANDPIPER
 - (ii) Finely streaked but not smudgy upper breast, grey-brown back, dark legs
 - (01) Bill less than head length SEMIPALMATED SANDPIPER
 - (02) Bill greater than 1.25X head length, drooped at tip, often reddish on back (NOTE: there is some overlap with Semipalmated, Western is accidental) WESTERN SANDPIPER







Shorebird Identification Tips

Notice the bird's...



Size — compare size to another bird in order to create a scale to learn where your bird fits among its bird group. Is it the size of a Mourning Dove, Robin, Red-winged Blackbird or Sparrow?

Leg Colour



Bill Length, Shape & Colour







Smaller than head Same length as head Longer than head





Curved Up

Curved Down

Plumage Colour & Pattern

Straight



Bars (horizontal lines) along body



Streaks on top of chest Streaks on top & sides Streaks along body





Bars along side



Eye Ring & Line







Distinct Eye Ring Indistinct Eye Ring No Eye Ring





Distinct Eye Line

Indistinct Eye Line No Eye Line

Tail & Wing Length



Wings longer than tail



Wings as long or shorter than tail



Teacher's Shorebird Answer Key

- 1. Black-bellied Plover
- 2. Piping Plover
- 3. Wilson's Phalarope (female)
- 4. Upland Sandpiper
- 5. Willet
- 6. Hudsonian Godwit
- 7. American Golden Plover
- 8. Killdeer (Plover)
- 9. Buff-breasted Sandpiper
- 10. Semipalmated Plover
- 11. American Woodcock
- 12. Short-billed Dowitcher
- 13. Marbled Godwit
- 14. Greater Yellowlegs
- 15. Spotted Sandpiper
- 16. Red-necked Phalarope
- 17. Long-billed Dowitcher
- 18. Wilson's Snipe
- 19. Pectoral Sandpiper
- 20. Solitary Sandpiper
- 21. Ruddy Turnstone
- 22. Stilt Sandpiper
- 23. Dunlin
- 24. White-rumped Sandpiper
- 25. Red Knot
- 26. Least Sandpiper
- 27. Baird's Sandpiper
- 28. Western Sandpiper
- 29. Semipalmated Sandpiper
- 30. Sanderling


Note:

When assigning a shorebird to your students, below are some tips regarding the difficulty level for each bird.

For students that may have the most difficulty in either focusing or detail distinction, it is recommend they are given a Plover (five in total; 1, 2, 7, 8, 10) to identity. Plovers have very distinct features and once students identify them as part of the Plover group, it significantly reduces the possible answers the students can choose.

Other shorebirds in this list that have very distinct features include the Dunlin (23), Ruddy Turnstone (21), Spotted Sandpiper (15), the Wilson's Phalarope (3), and the Red-necked Phalarope (16).

For students who are usually quick at tasks, detail-oriented and love a challenge, the most difficult shorebirds to identify (from this list) will be the Short-billed Dowitcher (12) and the Long-billed Dowitcher (17). However, these two species are difficult to distinguish so you may wish not to assign them if you think it would be too difficult for your students.


































Photo from Ducks Unlimited Canada







Photo by Francesco Veronesi, Flickr

Photo from Ducks Unlimited Canada





Smaller than a Mourning Dove, larger than a Red-winged Blackbird





Photo by Andrew Cannizzaro, Flickr








Photo by Omar Runolfsson, Flickr





Smaller than a Mourning Dove, larger than a Red-winged Blackbir Photo from Bettina Arrigoni, Wikipedia.







